

Hard exclusive electroproduction of two pions and their resonances

M.V. Polyakov

Petersburg Nuclear Physics Institute, Gatchina, St. Petersburg 188350, Russia
and
Institut für Theoretische Physik II, Ruhr-Universität Bochum, D-44780 Bochum,
Germany

Abstract

We study the hard exclusive production of two pions in the virtual photon fragmentation region with various invariant masses including the resonance region. The amplitude is expressed in terms of two-pion light cone distribution amplitudes (2π DA's). We derive dispersion relations for these amplitudes, which enables us to fix them completely in terms of $\pi\pi$ scattering phases and a few low-energy subtraction constants determined by the effective chiral lagrangian. Quantitative estimates of the resonance as well $\pi\pi$ background DA's at low normalization point are made. We also prove certain new soft pion theorem relating two-pion DA's to the one-pion DA. Crossing relations between 2π DA's and parton distributions in a pion are discussed.

We demonstrate that by studying the *shape* of the $\pi\pi$ mass spectra (not absolute cross section!) in a diffractive electroproduction one can extract the deviation of the meson (π, ρ , etc.) wave functions from their asymptotic form $6z(1-z)$ and hence to get important information about the structure of mesons. Also we discuss how the moments of quark distributions in a pion can be measured in the hard diffraction. We suggest (alternative to Söding's) parametrization of $\pi\pi$ spectra which is suitable for large photon virtuality.

Contents

1	Introduction	3
2	Two-pion light-cone distribution amplitudes.	4
3	Production of soft pions	6
4	Crossing relations between 2πDA's and quark distributions in a pion	8
5	Two-pion distribution amplitudes in the resonance region and DA's of resonances	11
5.1	Distribution amplitudes of resonances	13
6	Fixing low-energy constants from instantons	16
6.1	Isovector 2π DA's	17
6.2	Note on isoscalar 2π DA's	21
7	Two-pion distribution amplitudes in hard pions production processes	22
8	Conclusions	24
9	Acknowledgments	24
A	Appendix	25

1 Introduction

Recently it became possible to measure with high precision two pion hard diffractive production in the deeply virtual photon fragmentation region. The paper contains a detailed study of the “upper blob” of this process, more precisely, of the two-pion distribution amplitude.

It has been realized recently [1, 2, 3] that a number of hard exclusive processes of the type:

$$\gamma_L^*(q) + T(p) \rightarrow F(q') + T'(p') , \quad (1.1)$$

at large Q^2 with $t = (p - p')^2$, $x_{Bj} = Q^2/2(pq)$ and $q'^2 = M_F^2$, fixed are amenable to QCD description. The factorization theorem [1] asserts that the amplitude of the process (1.1) has the following form:

$$\sum_{i,j} \int_0^1 dz \int dx f_{i/T}^{T'}(x, x - x_{Bj}; t, \mu) H_{ij}(Q^2 x/x_{Bj}, Q^2, z, \mu) \Phi_j^F(z, \mu) \\ + \text{power-suppressed corrections}, \quad (1.2)$$

where $f_{i/T}^{T'}$ is $T \rightarrow T'$ skewed parton distribution [4, 5, 1, 2, 3] (for a review see [6]), $\Phi_j^F(z, \mu)$ is the distribution amplitude of the hadronic state F (not necessarily one particle state), and H_{ij} is a hard part computable in pQCD as series in $\alpha_s(Q^2)$. Here we shall study general properties of the distribution amplitudes $\Phi_j^F(z, \mu)$ when the final hadronic state F is a two pion state ($F = \pi\pi$).

The 2π DA's were introduced recently in [7] in the context of the QCD description of the process $\gamma^*\gamma \rightarrow 2\pi$, first model estimate of the corresponding DA at the two-pion invariant mass below the resonance region was done in [8]. In the present paper we discuss how to describe the resonance region. We shall see that calculations of the two-pion DA in low energy region [8] and the knowledge of $\pi\pi$ scattering phases are sufficient to determine 2π DA's in a wide range of two-pion invariant masses.

For definiteness we shall consider hard electroproduction of two pions off the nucleon:

$$\gamma_L^* + N \rightarrow 2\pi + T' , \quad (1.3)$$

where the final state T' can be $N, \Delta, \pi N$, etc. The usage of 2π DA's to describe the process (1.3) has several advantages comparing to the case of single resonance DA's (ρ, ρ' , etc.):

- The 2π DA's are the most general objects to describe the form of the $\pi\pi$ mass spectra in diffractive production of pions. We shall see that by studying the shape of the two-pion mass spectra (not the absolute cross section!) in diffractive production of two pions one can extract the information about the deviation of the meson (π, ρ) DA's from their asymptotic form, *i.e.* essentially non-perturbative information about the structure of mesons.

- The usage of 2π DA's is anyhow necessary in studies of channels with strong nonresonant background, for example, in the case of hard electroproduction of two neutral pions.
- By crossing symmetry the 2π DA's are related to skewed parton distribution (SPD's) in a pion (see section 4). Were 2π DA's measured one could get information about SPD's in a pion.
- From theoretical point of view 2π DA's are necessary to define DA's of unstable resonances (ρ, ρ', f_2, ρ_3 , etc.).

The plan of this paper is as follows. Section 2 contains definitions and general properties of two-pion distribution amplitudes. In Section 3 we prove soft pion theorems for the 2π DA's. These theorems relate the 2π DA's to the DA's of a single pion in a limit when one of the produced pions is soft. In section 4 we derive relations between 2π DA's and quark distributions (skewed and not) in a pion. These relations can be, in principle, used for measurements of quark distributions in a pion in hard exclusive reactions. Also they constrain the 2π DA's by already known parton distributions in a pion. Section 5 is devoted to derivation of the dispersion relations for the 2π DA's. These dispersion relations allow us to express the 2π DA's in a wide range of invariant two-pion masses in terms of (known) pion-pion phase shifts and a few low-energy subtraction constants. In subsection 5.1 we explain how to extract resonance DA's from 2π DA's. In section 6 we estimate the low-energy subtraction constants in the instanton model of the QCD vacuum, also results for ρ -meson DA's are presented. In section 7 we discuss the applications and implications of the 2π DA's to the hard electroproduction of pions. Conclusions and an outlook are given in section 8.

2 Two-pion light-cone distribution amplitudes.

We shall consider the following twist-2 chirally even and odd two-pion distribution amplitudes (2π DA's)

$$\begin{aligned} \Phi_{\parallel}^{ab}(z, \zeta, w^2) &= \frac{1}{4\pi} \int dx^- e^{-\frac{i}{2}zP^+x^-} \\ &\times \textit{out} \langle \pi^a(p_1) \pi^b(p_2) | \bar{\psi}(x) \hat{n} T \psi(0) | 0 \rangle \Big|_{x^+=x_{\perp}=0}, \end{aligned} \quad (2.1)$$

$$\begin{aligned} \Phi_{\perp}^{ab}(z, \zeta, w^2) &= \frac{i f_{2\pi}^{\perp}}{4\pi w^2} \int dx^- e^{-\frac{i}{2}zP^+x^-} \\ &\times \textit{out} \langle \pi^a(p_1) \pi^b(p_2) | \bar{\psi}(x) \sigma_{\mu\nu} n^{\mu} P^{\nu} T \psi(0) | 0 \rangle \Big|_{x^+=x_{\perp}=0}, \end{aligned} \quad (2.2)$$

Here, n is a light-like vector ($n^2 = 0$), which we choose to be $n_{\mu} = (1, 0, 0, 1)$. For any vector, V , the “plus” light-cone coordinate is defined as $V^+ \equiv (n \cdot V) = V^0 + V^3$; the “minus” component as $V^- = V^0 - V^3$. The outgoing pions have momenta p_1, p_2 , and $P \equiv p_1 + p_2$ is the total momentum of the final state, with $P^2 = w^2$. Finally, T is a

flavor matrix ($T = 1/2$ for the isosinglet, $T = \tau^3/2$ for the isovector 2π DA). [In the above equations the path-ordered exponential of the gauge field, required by gauge invariance is always assumed]. The constant of dimension of mass $f_{2\pi}^\perp$ is defined as matrix element of local operator:

$$\lim_{w^2 \rightarrow 0} \langle \pi^a(p_1) \pi^b(p_2) | \bar{\psi}(0) \sigma^{\mu\nu} \frac{\tau^3}{2} \psi(0) | 0 \rangle = \varepsilon^{ab3} \frac{2i}{f_{2\pi}^\perp} (p_1^\mu p_2^\nu - p_1^\nu p_2^\mu). \quad (2.3)$$

The generalized distribution amplitudes, eqs. (2.1,2.2), depend on the following kinematical variables: the quark momentum fraction with respect to the total momentum of the two-pion state, z ; the variable $\zeta \equiv p_1^+/P^+$ characterizing the distribution of longitudinal momentum between the two pions, and the invariant mass (c.m. energy), $w^2 = P^2$.

In what follows we shall work in the reference frame where $P_\perp = 0$. In this frame

$$P^- = \frac{w^2}{P^+}, \quad p_1^- = \frac{w^2(1-\zeta)}{P^+}, \quad p_\perp^2 \equiv p_{1\perp}^2 = p_{2\perp}^2 = w^2\zeta(1-\zeta) - m_\pi^2. \quad (2.4)$$

From these relations one obtains the following kinematical constraint:

$$\zeta(1-\zeta) \geq \frac{m_\pi^2}{w^2}. \quad (2.5)$$

The isospin decomposition of the 2π DA's eqs. (2.1,2.2) has the form¹:

$$\Phi^{ab} = \delta^{ab} \text{tr}(T) \Phi^{I=0} + \frac{1}{2} \text{tr}([\tau^a, \tau^b] T) \Phi^{I=1}. \quad (2.6)$$

From the C-parity one can easily derive the following symmetry properties of the isoscalar ($I = 0$) and isovector ($I = 1$) parts of 2π DA's eqs. (2.1,2.2):

$$\begin{aligned} \Phi^{I=0}(z, \zeta, w^2) &= -\Phi^{I=0}(1-z, \zeta, w^2) = \Phi^{I=0}(z, 1-\zeta, w^2), \\ \Phi^{I=1}(z, \zeta, w^2) &= \Phi^{I=1}(1-z, \zeta, w^2) = -\Phi^{I=1}(z, 1-\zeta, w^2). \end{aligned} \quad (2.7)$$

The isospin one parts of 2π DA's eqs. (2.1,2.2) are normalized as follows:

$$\begin{aligned} \int_0^1 dz \Phi_{\parallel}^{I=1}(z, \zeta, w^2) &= (2\zeta - 1) F_\pi(w^2), \\ \int_0^1 dz \Phi_{\perp}^{I=1}(z, \zeta, w^2) &= (2\zeta - 1) F_t(w^2), \end{aligned} \quad (2.8)$$

where $F_\pi(w^2)$ is the pion e.m. form factor in the time-like region ($F_\pi(0) = 1$) and $F_t(w^2)$ is the “tensor” form factor of the pion normalized by $F_t(0) = 1$.

Let us decompose 2π DA's in eigenfunctions of the ERBL [9] evolution equation (Gegenbauer polynomials $C_n^{3/2}(2z-1)$) and in partial waves of produced pions² (Gegenbauer

¹ We do not specify the indices ($\parallel$ and $\perp$) if an equation holds for both chirally even and odd DA's

² In the formulae below we, for simplicity, neglect threshold effects. How to take them into account is discussed in the Appendix.

polynomials $C_l^{1/2}(2\zeta - 1)$, see below). Generically the decomposition (for both isoscalar and isovector DA's) has the form:

$$\Phi(z, \zeta, w^2) = 6z(1 - z) \sum_{n=0}^{\infty} \sum_{l=0}^{n+1} B_{nl}(w^2) C_n^{3/2}(2z - 1) C_l^{1/2}(2\zeta - 1). \quad (2.9)$$

Using the symmetry properties (2.7) we see that the index n goes over even (odd) and index l goes over odd (even) numbers for isovector (isoscalar) 2π DA's. The normalization conditions (2.8) correspond to $B_{01}^{\parallel}(w^2) = F_{\pi}(w^2)$ and $B_{01}^{\perp}(0) = 1$.

The Gegenbauer polynomials $C_n^{3/2}(2z - 1)$ are eigenfunctions of the ERBL [9] evolution equation and hence the coefficients B_{nl} are renormalized multiplicatively (for even n and odd l):

$$B_{nl}(w^2; \mu) = B_{nl}(w^2; \mu_0) \left(\frac{\alpha_s(\mu)}{\alpha_s(\mu_0)} \right)^{(\gamma_n - \gamma_0)/(2\beta_0)}, \quad (2.10)$$

where $\beta_0 = 11 - 2/3N_f$ and the one loop anomalous dimensions are [10]:

$$\begin{aligned} \gamma_n^{\parallel} &= \frac{8}{3} \left(1 - \frac{2}{(1+n)(2+n)} + 4 \sum_{k=2}^{n+1} \frac{1}{k} \right), \\ \gamma_n^{\perp} &= \frac{8}{3} \left(1 + 4 \sum_{k=2}^{n+1} \frac{1}{k} \right). \end{aligned}$$

From the decomposition (2.9) and eq. (2.10) we can immediately make a simple prediction for the ratio of the hard P - and F - waves production amplitudes of pions in the reaction $\gamma^* p \rightarrow 2\pi p$ at large virtuality of the incident photon and fixed w^2 and Bjorken x :

$$\frac{F\text{-wave amplitude}}{P\text{-wave amplitude}} \sim \frac{1}{\log(Q^2)^{50/(99-6N_f)}}, \quad (2.11)$$

or generically for the $2k + 1$ wave:

$$\frac{(2k+1)\text{-wave amplitude}}{P\text{-wave amplitude}} \sim \frac{1}{\log(Q^2)^{\gamma_{2k}^{\parallel}/(2\beta_0)}}. \quad (2.12)$$

3 Production of soft pions

In this section we prove the low-energy theorem for the isovector $\Phi_{\parallel}^{I=1}(z, \zeta, w^2)$ relating this 2π DA to the single pion DA $\varphi_{\pi}(z)$ [11]. The latter is defined as:

$$\langle \pi^0(P) | \bar{\psi}(x) \hat{n} \gamma_5 \frac{\tau^3}{2} \psi(0) | 0 \rangle = i f_{\pi} P^+ \int_0^1 dz e^{izP \cdot x} \varphi_{\pi}(z). \quad (3.1)$$

Here $f_\pi = 93$ MeV is the pion decay constant.

Let us consider the matrix element ${}_{out}\langle \pi^a(p_1) \pi^b(p_2) | \bar{\psi}(x) \hat{n} \frac{\tau^3}{2} \psi(0) | 0 \rangle$ entering the definition of the chirally even isovector 2π DA eq. (2.1) and consider the kinematical limit when the 4-momentum of one of the pions goes to zero. This limit corresponds to $w^2 \rightarrow 0$ and $\zeta \rightarrow 1$ (or $\zeta \rightarrow 0$). Using the LSZ reduction formula we can write (for definiteness we take $p_2^\mu \rightarrow 0$ which corresponds to $w^2 \rightarrow 0$ and $\zeta \rightarrow 1$):

$$\begin{aligned} {}_{out}\langle \pi^a(p_1) \pi^b(p_2) | \bar{\psi}(x) \hat{n} \frac{\tau^3}{2} \psi(0) | 0 \rangle &= i \int d^4y e^{i(p_2 \cdot y)} (\partial_y^2 + m_\pi^2) \\ &\times \langle \pi^a(p_1) | T \left\{ \phi_\pi^b(y) \bar{\psi}(x) \hat{n} \frac{\tau^3}{2} \psi(0) \right\} | 0 \rangle, \end{aligned} \quad (3.2)$$

where $\phi_\pi^b(y)$ is a pion interpolating field which, owing to PCAC, can be related to the divergence of the axial current $\phi_\pi^b(y) = \frac{1}{f_\pi m_\pi^2} \partial_\mu J_5^{\mu b}(y)$. Substituting this into eq. (3.2) and taking the limit $p_2^\mu \rightarrow 0$ we can rewrite r.h.s. of eq. (3.2) in the form:

$$\begin{aligned} \frac{i}{f_\pi} \int d^4y \langle \pi^a(P) | T \left\{ \partial_\mu J_5^{\mu b}(y) \bar{\psi}(x) \hat{n} \frac{\tau^3}{2} \psi(0) \right\} | 0 \rangle &= \\ \frac{1}{i f_\pi} \int d^4y \left\{ \delta(y^0) \langle \pi^a(P) | \bar{\psi}(x) \hat{n} \frac{\tau^3}{2} [J_5^{0b}(y), \psi(0)] | 0 \rangle \right. & \quad (3.3) \\ \left. + \delta(y^0 - x^0) \langle \pi^a(P) | [J_5^{0b}(y), \bar{\psi}(x)] \hat{n} \frac{\tau^3}{2} \psi(0) | 0 \rangle \right\}. \end{aligned}$$

The equal time commutators entering the above equation are simply the global axial transformations of the quark fields:

$$\begin{aligned} \int d^4y \delta(y^0) [J_5^{0b}(y), \psi(0)] &= i \frac{\tau^b}{2} \gamma_5 \psi(0), \\ \int d^4y \delta(y^0 - x^0) [J_5^{0b}(y), \bar{\psi}(x)] &= i \bar{\psi}(x) \frac{\tau^b}{2} \gamma_5. \end{aligned} \quad (3.4)$$

Substituting this result into eq. (3.4) we get the following soft pion theorem:

$$\lim_{p_2 \rightarrow 0} {}_{out}\langle \pi^a(p_1) \pi^b(p_2) | \bar{\psi}(x) \hat{n} \frac{\tau^3}{2} \psi(0) | 0 \rangle = \frac{i \varepsilon^{3bc}}{f_\pi} \langle \pi^a(P) | \bar{\psi}(x) \hat{n} \gamma_5 \frac{\tau^c}{2} \psi(0) | 0 \rangle. \quad (3.5)$$

Applying this soft pion theorem to the matrix elements entering the definition of the chirally even 2π DA eq. (2.1) and comparing the resulting expression with definition of the pion DA, eq. (3.1), we get the following relation between the pion DA and the isovector chirally even 2π DA:

$$\Phi_{\parallel}^{I=1}(z, \zeta = 1, w^2 = 0) = -\Phi_{\parallel}^{I=1}(z, \zeta = 0, w^2 = 0) = \varphi_\pi(z). \quad (3.6)$$

The analogous theorem for the isoscalar part of the 2π DA's has the form:

$$\Phi_{\parallel}^{I=0}(z, \zeta = 1, w^2 = 0) = \Phi_{\parallel}^{I=0}(z, \zeta = 0, w^2 = 0) = 0. \quad (3.7)$$

Some comments are in order here. The derivation of the soft pion theorems presented here can be easily extended to other cases when we have an emission of soft pions. We should only keep in mind that for the case when we have nucleons in the *out* state (a typical example is a generalized skewed parton distributions, see [12, 13]) the singularities related to the emission of soft pions from the nucleon legs must be taken into account.

The soft pion theorem eq. (3.6) allows us to relate the coefficients $B_{nl}^{\parallel}(w^2)$ (see eq. (2.9)) and the coefficients in the expansion of the pion DA in the Gegenbauer polynomials

$$\varphi_{\pi}(z) = 6z(1-z) \left(1 + \sum_{n=\text{even}} a_n^{(\pi)} C_n^{3/2}(2z-1) \right). \quad (3.8)$$

The relation has the form:

$$a_n^{(\pi)} = \sum_{l=1}^{n+1} B_{nl}^{\parallel(I=1)}(0). \quad (3.9)$$

The soft pion theorem for the isoscalar chirally even 2π DA eq. (3.7) implies that

$$\sum_{l=0}^{n+1} B_{nl}^{\parallel(I=0)}(0) = 0. \quad (3.10)$$

4 Crossing relations between 2π DA's and quark distributions in a pion

An interesting feature of the pion is the fact that the 2π DA's are related to the skewed parton distribution in a pion by crossing. On one hand, the crossing relations derived in this section³ allow to use the information about quark distributions in a pion to constrain two-pion distribution amplitudes. On other hand, the measurements of 2π DA's in hard exclusive reactions can be used as a source of information on skewed quark distributions in a pion. Additionally the crossing relation are helpfull in modelling of skewed parton distributions.

The matrix element entering in the definition of 2π DA (2.1) by crossing is related to the matrix element in definition of skewed distribution:

$$\int \frac{d\lambda}{2\pi} \exp[i\lambda\tau n \cdot (p + p')] \langle \pi^a(p') | T \{ \bar{\psi}_{f'}(-\lambda n/2) \hat{n} \psi_f(\lambda n/2) \} | \pi^b(p) \rangle = \delta^{ab} \delta_{ff'} H^{I=0}(\tau, \xi, t) + i\epsilon^{abc} \tau_{ff'}^c H^{I=1}(\tau, \xi, t), \quad (4.1)$$

where ξ is a skewedness parameter defined as: $\xi = -(n \cdot (p' - p)) / (n \cdot (p' + p))$, $t = (p' - p)^2$ and light-cone vector n^μ introduced in section 2.

³Part of the material of this section was presented already in recent preprints [14, 15].

This allows to express the moments of skewed parton distributions in terms of expansion coefficients B_{nl} in eq. (2.9):

$$\int_{-1}^1 d\tau \tau^{N-1} H^I(\tau, \xi, t) = \sum_{n=0}^{N-1} \sum_{l=0}^{n+1} B_{nl}^I(t) \xi^N C_l^{1/2}\left(\frac{1}{\xi}\right) \int_{-1}^1 dx \frac{3}{4} [1-x^2] x^{N-1} C_n^{3/2}(x). \quad (4.2)$$

Let us note that this expression satisfy the polynomiality condition for moments of SPD's (see ref. [6]).

In order to prove the relation (4.2) let us consider the operator expression for the N -th moments of SPD. It is given by off-forward matrix element of local operator of spin N and twist-2:

$$\int_{-1}^1 d\tau \tau^{N-1} H(\tau, \xi, t) = \frac{1}{[(p' + p)^+]^N} \langle p' | \bar{\psi} \gamma^+ (\overleftrightarrow{\nabla}^+)^{N-1} \psi | p \rangle, \quad (4.3)$$

Analogously the N -th moments of 2π DA (2.1) is expressed in terms of the following matrix element of the same local operator:

$$\begin{aligned} \int_0^1 dz (2z-1)^{N-1} \Phi^I(z, \zeta, w^2) &= \frac{1}{[(p' + p)^+]^N} \langle p' p | \bar{\psi} \gamma^+ (\overleftrightarrow{\nabla}^+)^{N-1} \psi | 0 \rangle = \\ &= \sum_{n=0}^{N-1} \sum_{l=0}^{n+1} B_{nl}^I((p' + p)^2) C_l^{1/2}\left(\frac{(p' - p)^+}{(p' + p)^+}\right) \int_{-1}^1 dx \frac{3}{4} [1-x^2] x^{N-1} C_n^{3/2}(x), \end{aligned} \quad (4.4)$$

where in the last equality we used the double expansion (2.9). The matrix elements in eq. (4.4) and in eq. (4.3) are related to each other by crossing:

$$\begin{aligned} \frac{1}{[(p + p')^+]^N} \langle p' | \bar{\psi} \gamma^+ (\overleftrightarrow{\nabla}^+)^{N-1} \psi | p \rangle &= \frac{1}{[(p' + p)^+]^N} \langle -p, p' | \bar{\psi} \gamma^+ (\overleftrightarrow{\nabla}^+)^{N-1} \psi | 0 \rangle = \\ \frac{[(p' - p)^+]^N}{[(p' + p)^+]^N} \sum_{n=0}^{N-1} \sum_{l=0}^{n+1} B_{nl}^I((p' - p)^2) C_l^{1/2}\left(\frac{(p' + p)^+}{(p' - p)^+}\right) \int_{-1}^1 dx \frac{3}{4} [1-x^2] x^{N-1} C_n^{3/2}(x) &= \\ \sum_{n=0}^{N-1} \sum_{l=0}^{n+1} B_{nl}^I(t) \xi^N C_l^{1/2}\left(\frac{1}{\xi}\right) \int_{-1}^1 dx \frac{3}{4} [1-x^2] x^{N-1} C_n^{3/2}(x). \end{aligned} \quad (4.5)$$

This is exactly relation (4.2). Let us note that in the expression (4.2) the coefficients $B_{nl}(t)$ enter at negative argument whereas in (2.9) at positive argument. The corresponding analytical continuation can be done with help of dispersion relations (5.8) (see the next section).

If we take the forward limit in (4.2), we obtain the relations between moments of quark distributions in a pion and coefficients $B_{nl}(0)$:

$$\begin{aligned} M_N^{(\pi)} &\equiv \int_0^1 dx x^{N-1} (q_\pi(x) - \bar{q}_\pi(x)) = B_{N-1,N}^{I=1}(0) A_N \quad \text{for odd } N, \\ M_N^{(\pi)} &\equiv 2 \int_0^1 dx x^{N-1} (q_\pi(x) + \bar{q}_\pi(x)) = B_{N-1,N}^{I=0}(0) A_N \quad \text{for even } N, \end{aligned} \quad (4.6)$$

where A_N are numerical coefficients (*e.g.* $A_1 = 1, A_2 = 9/5, A_3 = 6/7, A_4 = 5/3$, etc.) and $q_\pi(x) = u^{\pi^+}(x)$. For example, from eq. (4.6) we obtain that $B_{01}^{I=1}(0) = M_1^{(\pi)} = 1$ what corresponds to normalization condition (2.8). Also it is easy to see that $B_{12}^{I=0}(0) = 5/9 M_2^{(\pi)}$ corresponds to normalization for isoscalar 2π DA ⁴:

$$\int_0^1 dz (2z - 1) \Phi^{I=0}(z, \zeta, w^2 = 0) = -2 M_2^\pi \zeta (1 - \zeta) , \quad (4.7)$$

where M_2^π is a momentum fraction carried by quarks in a pion.

Using relations (4.6) and soft pion theorem (3.6) we can express the coefficient $B_{21}(0)$ [it describes the deviation of isovector 2π DA from its asymptotic form $\Phi_{as}^{I=1}(z, \zeta, w^2) = 6z(1-z)(2\zeta-1)F_\pi^{\text{e.m.}}(w^2)$] as:

$$B_{21}^{I=1}(0) = a_2^{(\pi)} - \frac{7}{6} M_3^{(\pi)} . \quad (4.8)$$

From the expression for the moments of SPD (4.2) we can formally construct the SPD itself as series:

$$H^I(\tau, \xi, t) = \sum_{n=0}^{\infty} \sum_{l=0}^{n+1} \frac{3}{4} \left(1 - \frac{\tau^2}{\xi^2}\right) \theta(|x| - |\xi|) B_{nl}^I(t) C_n^{3/2}\left(\frac{\tau}{\xi}\right) C_l^{1/2}\left(\frac{1}{\xi}\right) , \quad (4.9)$$

where each term in the sum has a support $-\xi < \tau < \xi$. Let us note however that it does not imply that the SPD $H(\tau, \xi, t)$ has the same support because generically the sum (4.9) is divergent at fixed τ and ξ . In the formal solution (4.9) each individual term of the series gives the contribution to the amplitude which behaves as $1/\xi^l$ at small ξ what would imply the violation of unitarity for the hard exclusive reactions. This “disaster” is avoided through the fact the series is divergent and one can not calculate the asymptotic behaviour term by term. This situation is similar to duality idea: the *entire* skewed and usual quark distributions are given by infinite sum over t -channel resonances. Of course, there might be “non-dual” or “two component” way to compensate the bad behaviour of the resonances by some equally bad behaviour of non-resonant background. Such “two component” model can be written as:

$$\begin{aligned} H^I(\tau, \xi, t) &= \sum_{l=0}^{L_0} \sum_{n=l+1}^{\infty} \frac{3}{4} \left(1 - \frac{\tau^2}{\xi^2}\right) \theta(|x| - |\xi|) B_{nl}^I(t) C_n^{3/2}\left(\frac{\tau}{\xi}\right) C_l^{1/2}\left(\frac{1}{\xi}\right) \\ &+ H_{\text{non-resonant}}^I(\tau, \xi, t) , \end{aligned} \quad (4.10)$$

where L_0 is the maximal spin of resonances which are taken into account and $H_{\text{non-resonant}}^I(\tau, \xi, t)$ is a non-resonant background. The latter can be modelled in terms of double distributions [16], however if $L_0 > 0$ the non-resonant contribution has to be oscillating function in order to avoid double counting in calculations of low moments of SPD ($N \leq L_0$) which are saturated by resonances in t -channel.

⁴To see this one has to use additionally soft pion theorem (3.7) for isoscalar 2π DA

To summarize, the representation of SPD's (4.9) in terms of only resonance exchanges gives a new picture of quark distributions in a pion which is *dual* to the picture based on the concept of double distributions [16]. It would be extremely interesting to build a model where the duality relation

$$\lim_{t, \xi \rightarrow 0} \sum_{n=0}^{\infty} \sum_{l=0}^{n+1} \frac{3}{4} (1 - \frac{\tau^2}{\xi^2}) \theta(|x| - |\xi|) B_{nl}^I(t) C_n^{3/2}(\frac{\tau}{\xi}) C_l^{1/2}(\frac{1}{\xi}) = q_{\pi}(\tau) , \quad (4.11)$$

is realized explicitly.

At the end let us note that from the formal solution (4.9) and soft pion theorems (3.6, 3.7) we can determine the shape of SPD's in a pion in a limiting case: $\xi = 1$ and $t = 0$:

$$H^{I=1}(\tau, \xi = 1, t = 0) = \frac{1}{2} \varphi_{\pi}(\frac{\tau + 1}{2}) , \quad H^{I=0}(\tau, \xi = 1, t = 0) = 0 ,$$

where $\varphi_{\pi}(z)$ is a pion distribution amplitude. These limiting relations impose strong constraints on the models of skewed parton distributions. For example it is easy to see that the model based on double distributions proposed in [16] does not satisfy these constraints.

5 Two-pion distribution amplitudes in the resonance region and DA's of resonances

Generally speaking, the 2π DA's are complex functions owing to the strong interaction of the produced pions. Above the two-pion threshold $w^2 = 4m_{\pi}^2$ the 2π DA's develop an imaginary part corresponding to the contribution of on-shell intermediate states (2π , 4π , etc.). In the region $w^2 < 16m_{\pi}^2$ the imaginary part is related to the pion-pion scattering amplitude by the Watson theorem [17]. This relation can be written in the following form (in this section we omit the indices $\parallel$ and $\perp$ but show the isospin index, since all formulae in this section apply to both chirally even and odd 2π DA's):

$$\text{Im} \Phi^I(z, \zeta, w^2) = \int d(\text{phase space}) \Phi^I(z, \zeta', w^2)^* A_{\pi\pi}^I(\zeta', \vec{p}'_{\perp} | \zeta, \vec{p}_{\perp}) , \quad (5.1)$$

where $A_{\pi\pi}^I(\zeta', \vec{p}'_{\perp} | \zeta, \vec{p}_{\perp})$ is pion-pion scattering amplitude with isospin I in the s -channel, ζ' and $p'_{\perp}$ are the momenta of the intermediate pion, the phase space integral in these variables has the form:

$$\int d(\text{phase space}) = \frac{1}{64\pi^2} \int_0^1 d\zeta' \int_0^{2\pi} d\theta_{\perp} , \quad (5.2)$$

where $\theta_{\perp}$ is the angle between $p_{\perp}$ and $p'_{\perp}$. In order to find the form of the $\pi\pi$ scattering amplitude in the light-cone variables we start with the usual partial wave expansion of the pion-pion scattering amplitude:

$$A_{\pi\pi}^I = 32\pi \sum_l (2l+1) \sin[\delta_l^I(w^2)] e^{i\delta_l^I(w^2)} P_l[\cos(\theta_{cm})] , \quad (5.3)$$

where θ_{cm} is the scattering angle in the centre of mass system, it can be easily expressed in terms of light-cone variables. The relation has the form (here for a massless pion for simplicity)⁵:

$$\cos(\theta_{cm}) = 1 - 2(\zeta + \zeta' - 2\zeta\zeta') - 4\sqrt{\zeta(1-\zeta)\zeta'(1-\zeta')} \cos(\theta_\perp) . \quad (5.4)$$

Substituting the partial wave expansion (5.3) into eq. (5.1) and computing the integral over $\theta_\perp$ using the identity:

$$\int_0^{2\pi} d\theta_\perp P_l \left[1 - 2(\zeta + \zeta' - 2\zeta\zeta') - 4\sqrt{\zeta(1-\zeta)\zeta'(1-\zeta')} \cos(\theta_\perp) \right] = 2\pi C_l^{1/2}(2\zeta-1)C_l^{1/2}(2\zeta'-1) , \quad (5.5)$$

we get the following expression for the imaginary part of the 2π DA:

$$\begin{aligned} \text{Im } \Phi^I(z, \zeta, w^2 + i0) &= \sum_l (2l+1) \int_0^1 d\zeta' \Phi^I(z, \zeta', w^2)^* \sin[\delta_l^I(w^2)] e^{i\delta_l^I(w^2)} \\ &\times C_l^{1/2}(2\zeta-1)C_l^{1/2}(2\zeta'-1) . \end{aligned} \quad (5.6)$$

If we now substitute the expansion of the 2π DA in the Gegenbauer polynomials (2.9) in eq. (5.6) we get the equation for the imaginary part of the expansion coefficients B_{nl} :

$$\text{Im } B_{nl}^I(w^2) = \sin[\delta_l^I(w^2)] e^{i\delta_l^I(w^2)} B_{nl}^I(w^2)^* = \tan[\delta_l^I(w^2)] \text{Re } B_{nl}^I(w^2) . \quad (5.7)$$

One sees immediately that the phases of the coefficients B_{nl}^I are the same as the phase shifts δ_l^I . Let us also note that the equation (5.7) for the case $n=0, l=1$ and $I=1$ is identical to that obtained in [18] for the pion electromagnetic form factor, which is not surprising because of identity $B_{01}(w^2) = F_\pi(w^2)$. Since the form of equation (5.7) is very similar to that for e.m. pion form factor one can analyze it following ref. [18].

Using eq. (5.7) one can write an N -subtracted dispersion relation for the $B_{nl}^I(w^2)$:

$$B_{nl}^I(w^2) = \sum_{k=0}^{N-1} \frac{w^{2k}}{k!} \frac{d^k}{dw^{2k}} B_{nl}^I(0) + \frac{w^{2N}}{\pi} \int_{4m_\pi^2}^\infty ds \frac{\tan \delta_l^I(s) \text{Re } B_{nl}^I(s)}{s^N (s - w^2 - i0)} . \quad (5.8)$$

The solution of such type of dispersion relation was found long ago by Omnes [19] and has the exponential form:

⁵Expressions for massive case see in Appendix.

$$B_{nl}^I(w^2) = B_{nl}^I(0) \exp \left\{ \sum_{k=1}^{N-1} \frac{w^{2k}}{k!} \frac{d^k}{dw^{2k}} \log B_{nl}^I(0) + \frac{w^{2N}}{\pi} \int_{4m_\pi^2}^{\infty} ds \frac{\delta_l^I(s)}{s^N (s - w^2 - i0)} \right\}. \quad (5.9)$$

This equation, by derivation, is valid below the inelastic threshold $w^2 < 16m_\pi^2$, however the inelasticities (at least for the isospin one channel) in the pion-pion scattering are small up to rather large energies. Also the equation (5.9) being applied to the case of the pion form factor ($n = 0, l = 1$) [18, 20] gives an excellent description of experimental data up to $w^2 \approx 2.5 \text{ GeV}^2$ with two subtractions which are fixed by the value of the pion e.m. form factor at zero and its charge radius (see figure in [20]). Moreover the solution (5.9) can be systematically improved by taking into account higher thresholds (see discussion in [21]), probably the most important is the $K\bar{K}$ one. The contribution of higher intermediate states can be suppressed by choosing sufficiently large number of subtractions N , see discussion in section 5.1.

A great advantage of the solution eq. (5.9) is that it gives the 2π DA's in a wide range of energies in terms of known $\pi\pi$ phase shifts and a few subtraction constants (usually two is sufficient). The key observation is that these constants are related to the *low-energy* behaviour at $w^2 \rightarrow 0$ of the 2π DA's. In the low energy region they can be computed using the effective chiral quark-pion lagrangian. In the next section we shall compute the subtraction constants using the low-energy effective lagrangian derived from the instanton model of QCD vacuum [22]. For the first estimates of the 2π DA in this model in low energy region see [8]. After the subtraction constants are fixed (see section 6) we shall know the 2π DA's in a broad region of invariant masses. Of particular interest is the case when w^2 is close to the mass of some resonance (say ρ): in this case we shall be able to extract DA of this resonance from the solution (5.9), this is explained in the next subsection.

5.1 Distribution amplitudes of resonances

The light-cone distribution amplitudes of mesons are defined as matrix elements of quark-antiquark non-local operators at light-like separations [23]. Here we consider as an example the ρ -mesons DA's, all equations in this section can be easily generalized to other mesons. Specifically, we consider chirally even and odd leading twist ρ -meson DA's defined as:

$$\langle \rho^0(P) | \bar{\psi}(x) \hat{n} \frac{\tau^3}{2} \psi(0) | 0 \rangle \Big|_{x^+ = x_\perp = 0} = (n \cdot e^{(\lambda)}) \frac{f_\rho^\parallel m_\rho}{\sqrt{2}} \int_0^1 dz e^{-iz(P \cdot x)} \varphi_\rho^\parallel(z), \quad (5.10)$$

$$\langle \rho^0(P) | \bar{\psi}(x) \sigma_{\mu\nu} n^\mu P^\nu \frac{\tau^3}{2} \psi(0) | 0 \rangle \Big|_{x^+ = x_\perp = 0} = i(n \cdot e^{(\lambda)}) \frac{f_\rho^\perp m_\rho^2}{\sqrt{2}} \int_0^1 dz e^{iz(P \cdot x)} \varphi_\rho^\perp(z), \quad (5.11)$$

where $e_\mu^{(\lambda)}$ is the polarization vector of the ρ -meson. The normalization constants $f_\rho^\perp$ and $f_\rho^\parallel$ have dimension of mass and are chosen in such a way that (below we suppress indices $\perp$ and $\parallel$ where it does not lead to confusion):

$$\int_0^1 dz \varphi_\rho(z) = 1 .$$

The light-cone ρ meson DA's are used to describe hard leptoproduction of vector mesons [24], and in applications of light-cone QCD sum rules for exclusive semileptonic and radiative weak decays of the B -meson [25].

There are certain problems with the definitions eqs. (5.10,5.11), because the ρ -meson is an unstable particle. One can treat the ρ -meson as a quasistable particle, *i.e.* consider the formal limit $\Gamma_\rho \rightarrow 0$. From experimental point of view ρ -meson is detected as a Breit-Wigner peak in the invariant mass distribution of produced pions. Therefore a more adequate (and accurate) way to describe the hard production experiments is in the terms of two-pion DA's.

In order to extract the rho meson DA's from the 2π DA's we use the expression (5.9) of 2π DA in terms of the pion-pion phase shifts. In the vicinity of the resonance the corresponding phase shift (we shall speak for definiteness about ρ -meson, *i.e.* $l = 1$, isospin one) can be approximated by the Breit-Wigner formula:

$$\delta_1^1(w^2 \sim m_\rho^2) = \arctan\left(\frac{m_\rho \Gamma_\rho}{m_\rho^2 - w^2}\right) . \quad (5.12)$$

In the limit of vanishing width the contribution of the resonance part of the phase shift in the integral in eq. (5.9) can be written as :

$$\begin{aligned} \lim_{\Gamma_\rho \rightarrow 0} \frac{w^{2N}}{\pi} \int_{4m_\pi^2}^\infty ds \frac{\delta_1^1(s)}{s^N(s - w^2 - i0)} &= i\frac{\pi}{2} - \log(m_\rho^2 - w^2) + \log m_\rho^2 \\ &+ (\text{pol. of order } N - 1) . \end{aligned} \quad (5.13)$$

This result being exponentiated gives rise to the meson pole. Matching the solution for 2π DA's eq. (5.9) near $w^2 \sim m_\rho^2$ to the Breit-Wigner form we get the a relation between ρ meson DA's and 2π DA's. We obtain the following expression for the coefficients of the expansion of ρ meson DA'S in Gegenbauer polynomials

$$\varphi_\rho(z) = 6z(1-z) \left(1 + \sum_{n=\text{even}} a_n^{(\rho)} C_n^{3/2}(2z-1) \right),$$

in terms of subtraction constants entering eq. (5.9) only

$$a_n^{(\rho)} = B_{n1}(0) \exp\left(\sum_{k=1}^{N-1} c_k^{(n1)} m_\rho^{2k}\right) , \quad (5.14)$$

where the subtraction constants $c_k^{(nl)}$ can be expressed in terms of $B_{nl}(w^2)$ at low energies

$$c_k^{(nl)} = \frac{1}{k!} \frac{d^k}{dw^{2k}} [\log B_{nl}(w^2) - \log B_{l-1l}(w^2)] \Big|_{w^2=0} . \quad (5.15)$$

The constants $f_\rho^{\parallel,\perp}$ can be computed as:

$$\begin{aligned} f_\rho^\parallel &= \frac{\sqrt{2}\Gamma_\rho \text{Im} B_{01}^\parallel(m_\rho^2)}{g_{\rho\pi\pi}}, \\ f_\rho^\perp &= \frac{\sqrt{2}\Gamma_\rho m_\rho \text{Im} B_{01}^\perp(m_\rho^2)}{g_{\rho\pi\pi} f_{2\pi}^\perp}, \end{aligned} \quad (5.16)$$

where $g_{\rho\pi\pi}$ is a ρ meson decay constant into two pions, $f_{2\pi}^\perp$ is defined by eq. (2.3). The constants $B_{01}^{\parallel,\perp}(m_\rho^2)$ are purely imaginary and can be computed using eq. (5.9) in terms of the pion-pion phase shift δ_1^1 and low-energy subtraction constants. The latter are actually pion charge radius and radius of pion tensor form factor (they are estimated in the next section).

eqs. (5.14,5.16) can be easily generalized to the DA's of higher spin resonances. For example, the chirally even DA's of an isovector resonance of spin j can be computed in terms of the 2π DA's as:

$$\phi_j(z) = 6z(1-z) \left(C_{j-1}^{3/2}(2z-1) + \sum_{n=j+1}^{\infty} a_n^{(j)} C_n^{3/2}(2z-1) \right), \quad (5.17)$$

with coefficients $a_n^{(j)}$ computed as:

$$a_n^{(j)} = \frac{B_{nj}(0)}{B_{j-1j}(0)} \exp\left(\sum_{k=1}^{N-1} c_k^{(nj)} m_R^{2k}\right). \quad (5.18)$$

The corresponding normalization constant is:

$$f_R = \frac{\sqrt{2}\Gamma_R \text{Im} B_{j-1j}(m_R^2)}{g_{R\pi\pi}}. \quad (5.19)$$

The scale dependence of f_R and $a_n^{(j)}$ is obvious.

We see that the ρ meson DA's (we speak for definiteness about ρ meson, but all arguments below apply also to any other resonance) are fixed in terms of the subtraction (low-energy) constants, which would be all fixed if we knew the exact low-energy lagrangian of QCD⁶. The latter is not fully known, so we shall resort to the low-energy lagrangian derived from the instanton model of the QCD vacuum [22].

The r.h.s. of eq. (5.14) depends on the number of subtractions N whereas its l.h.s. is independent of this number. At first glance this fact implies that the subtraction constants $c_k^{(nl)}$ are all zero. However we should keep in mind that the dispersion relation eq. (5.8) is not exact, in its derivation the contribution of intermediate states other than two pion was neglected. In order to suppress the contribution of the higher states we have to take

⁶For example, the constant $c_1^{\parallel(01)}$ is proportional to the parameter L_9 of Gasser-Leutwyler parametrization of the EChL [30].

a large enough number of subtractions in the dispersion relation (5.8). This implies that one can expect an (approximate) independence of the r.h.s. of eq. (5.14) of N only for sufficiently large N .

Increasing the number of subtractions suppresses the high energy region, where the approximations were made, and hence increases the accuracy of the approximate solution (5.9). However, increasing the number of subtractions we emphasize the low energy region the quantitative description of which is limited by incomplete knowledge of the effective chiral lagrangian. Therefore for phenomenological applications one has to choose some optimal N . The analysis of the pion e.m. form factor shows [18, 20] that already for $N = 2$ the contribution of the high energy states is sufficiently suppressed to describe the pion form factor with good accuracy up to $w^2 \approx 2.5 \text{ GeV}^2$.

In the next section we take $N = 2$. Since with this value the contribution of the high energy states in dispersion relation (5.8) is sufficiently well damped and simultaneously we can reliably use the approximate low energy quark-pion chiral lagrangian derived from the instanton model of QCD vacuum [22]. In language of the chiral perturbation theory $N = 2$ corresponds to the fixing of the effective chiral lagrangian (EChL) to the fourth order. To this order the effective chiral theory derived from the instanton model gives [22, 26] the constants of the fourth order EChL (actually only its part describing pions interacting with external vector and axial currents) which are very close to ones obtained in the phenomenological analysis of Gasser and Leutwyler [30].

6 Fixing low-energy constants from instantons

The technique how to compute matrix elements of bilocal quark operators at low energies in the effective low-energy quark-pion theory was explained in details in refs. [27, 28] (for pion and photon DA's) and in [8] (for 2π DA's). Here we only sketch the main points of the calculations, for details the reader should consult refs. [27, 28, 8].

We shall compute the 2π DA's at small w^2 , eq. (2.1,2.2), using the effective low-energy theory of pions interacting with massive “constituent” quarks, which has been derived from the instanton model of the QCD vacuum [22]. The coupling of the pion field to the quarks, whose form is restricted by the chiral invariance, is described by the action

$$S_{\text{int}} = \int d^4x \bar{\psi}(x) \sqrt{M(\partial^2)} U^{\gamma_5}(x) \sqrt{M(\partial^2)} \psi(x), \quad (6.1)$$

where

$$U^{\gamma_5}(x) = e^{i\gamma_5 \tau^a \pi^a(x)/f_\pi} = 1 + \frac{i}{f_\pi} \gamma_5 \tau^a \pi^a(x) - \frac{1}{2f_\pi^2} \pi^a(x) \pi^a(x) + \dots \quad (6.2)$$

Here, $f_\pi = 93 \text{ MeV}$ is the weak pion decay constant

The momentum-dependent dynamical quark mass, $M(p)$, plays the role of an UV regulator. Its form for Euclidean momenta was derived in ref.[22]. The momentum dependent mass cuts loop integrals at momenta of order of the inverse average instanton size, $\bar{\rho}^{-1} \approx 600 \text{ MeV}$. One should note that the value of the mass at zero momentum, $M(0)$, is parametrically small compared to $\bar{\rho}^{-1}$; the product $M(0)\bar{\rho}$ is proportional to the packing fraction of the instantons in the medium, which is a small algebraic parameter

fundamental to this picture. Numerically, a value $M(0) = 350 \text{ MeV}$ was obtained in ref. [22].

6.1 Isovector $2\pi\text{DA}$'s

Using the technique described in [8] we can easily see that the loop integral in the effective theory for the chirally odd isovector $2\pi\text{DA}$ (2.2) contains no divergencies, and hence we can neglect during its computation the momentum dependence of the constituent quark mass. Chirally even isovector $2\pi\text{DA}$ (2.1) was computed in [8], the computation there was done neglecting the momentum dependence of the constituent quark mass and absorbing the divergencies into f_π . It is easy to see that all (logarithmic) divergencies are contained in the w^2 -independent piece of this DA, the w^2 dependent piece is finite. The result of [8] in this approximation for $w^2 = 0$ was very simple:

$$\Phi_{\parallel}(z, \zeta, w^2 = 0) = \text{sign}(1 - 2\zeta)\theta[(z - \zeta)(1 - \zeta - z)] , \quad (6.3)$$

where $\theta(x)$ is a step function. We see that chirally even isovector $2\pi\text{DA}$ exhibits jumps at points $z = \zeta$ and $z = 1 - \zeta$. As was shown in [37, 27, 28] the momentum dependence of the constituent quark mass becomes crucial at these points. Here for the numerical estimates of the w^2 independent piece we employ a simple numerical fit to the momentum dependent constituent quark mass obtained from the instanton model of the QCD vacuum,

$$M(-p^2) = \frac{M_0}{(1 + 0.5 p^2 \bar{\rho}^2)^2} . \quad (6.4)$$

As we already mentioned the w^2 -dependent piece of the chirally even isovector $2\pi\text{DA}$ is finite (it does not exhibit the jumps) and hence it can be computed neglecting the UV-cutoff provided by the momentum dependence of the constituent quark mass.

Results of a computation of the chirally even isovector $2\pi\text{DA}$ at $w^2 = 0$ are shown in Fig. 1, where we have plotted $\Phi_{\parallel}^{I=1}(z, \zeta, w^2 = 0)$ as a function of z at various values of ζ . We see that the function is concentrated in the interval of z between ζ and $1 - \zeta$ and the jumps are smoothed. Also it is interesting to note that if we take $\zeta = 1$ the result for $\Phi_{\parallel}^{I=1}(z, \zeta, w^2 = 0)$ coincides exactly with that for the pion DA obtained in [27, 28] in the same model, what is not surprising due to the soft pion theorem (3.6).

Inspecting loop integrals for the chirally even $2\pi\text{DA}$'s in the model (see typical expressions in ref. [8]) we can make interesting observation: in the limits $\zeta \rightarrow 1$ and $m_\pi \rightarrow 0$ the w^2 dependence appear exclusively in the combination $z(1 - z)w^2$. This observation allows us to obtain the following model relations for the coefficients of Gegenbauer expansion (2.9) $B_{nl}^{\parallel}(w^2)$ (we omit the superscript $\parallel$ below to make the expression not so busy):

$$\sum_{\substack{l=1 \\ \text{odd}}}^{n+1} \frac{d^j}{dw^{2j}} B_{nl}^{I=1}(w^2)|_{w^2=0} = 0 , \quad \text{for any even } n \geq 2j , \quad (6.5)$$

$$\sum_{\substack{l=0 \\ \text{even}}}^{n+1} \frac{d^j}{dw^{2j}} B_{nl}^{I=0}(w^2)|_{w^2=0} = 0 , \quad \text{for any odd } n \geq 2j - 1 . \quad (6.6)$$

These *model* results, in a sense, can be viewed as a generalization of the soft pion theorems (3.9, 3.10). Although we find these relations in the model calculations, we can speculate that they have more general nature and follow from the chiral Ward identities in the limit of large colour number $N_c \rightarrow \infty$. This conjecture will be worked out in details elsewhere [36].

Below we give the results of numerical calculations of a few first coefficients of Gegenbauer expansion (2.9) $B_{nl}^{\parallel}(w^2)$ of the isovector chirally even 2π DA at low w^2 :

$$B_{01}^{\parallel}(w^2) = 1 + \frac{N_c w^2}{24\pi^2 f_\pi^2} + \dots, \quad (6.7)$$

$$B_{21}^{\parallel}(w^2) = -0.2 - \frac{7N_c w^2}{1440\pi^2 f_\pi^2} + \dots, \quad (6.8)$$

$$B_{23}^{\parallel}(w^2) = 0.26 + \frac{7N_c w^2}{1440\pi^2 f_\pi^2} + \dots, \quad (6.9)$$

$$B_{41}^{\parallel}(w^2) = -0.006 - \frac{11N_c w^2}{25200\pi^2 f_\pi^2} + \dots, \quad (6.10)$$

$$B_{43}^{\parallel}(w^2) = -0.12 - \frac{11N_c w^2}{10800\pi^2 f_\pi^2} + \dots, \quad (6.11)$$

$$B_{45}^{\parallel}(w^2) = 0.14 + \frac{11N_c w^2}{7560\pi^2 f_\pi^2} + \dots. \quad (6.12)$$

From this calculation we immediately can extract the coefficients of the Gegenbauer expansion of the pion DA (3.8) using the soft pion theorem (3.9):

$$a_2^{(\pi)} = 0.06, \quad a_4^{(\pi)} = 0.014. \quad (6.13)$$

These are exactly the values obtained in [28].

Remembering that $B_{01}^{\parallel}(w^2) = F_\pi(w^2)$ we extract from eq. (6.7) the value of the pion electromagnetic charge radius (without chiral loops!)

$$\langle r^2 \rangle_{\text{e.m.}} = \frac{N_c}{4\pi^2 f_\pi^2}, \quad (6.14)$$

which coincides with the result obtained previously in ref. [22]. The model result (6.14) numerically gives $\langle r^2 \rangle_{\text{e.m.}} = 0.35 \text{ fm}^2$ to be compared with the experimental value $0.439 \pm 0.008 \text{ fm}^2$ [29]. Comparing these numbers we should keep in mind that the model calculations do not include the chiral logarithms, the latter appear when we perform the integral in eq. (5.9) (for $n = 0, l = 1$) with the pion-pion phase shift obtained in the leading order of the chiral perturbation theory:

$$\delta_1^1(w^2) = \frac{w^2(1 - 4m_\pi^2/w^2)^{3/2}}{96\pi f_\pi^2}.$$

Then the result for the pion charge radius becomes (remember that $B_{01}(w^2) = F_\pi(w^2)$):

$$\langle r^2 \rangle_{\text{e.m.}} = \frac{N_c}{4\pi^2 f_\pi^2} - \frac{1}{16\pi^2 f_\pi^2} \log\left(\frac{m_\pi^2}{m_\rho^2}\right) \approx 0.45 \text{ fm}^2,$$

in good agreement with the experimental value.

Basing on the results (6.7-6.12) we can easily estimate the low-energy subtraction constants, see eqs. (5.9,5.15), the results are summarized in Table 1, they refer to the normalization point $\mu \sim 1/\bar{\rho} \approx 600 \text{ MeV}$ the inverse instanton size in the instanton vacuum.

Having the values of the low-energy subtraction constants we can obtain the chirally even isovector $2\pi\text{DA}$ in a wide interval of invariant two-pion masses using the solution (5.9) with $N = 2$ (see the discussion in previous section). The first qualitative observation we can make on basis of Table 1 is that the coefficients in front of $C_n^{3/2}(2z-1)$ for $n = 2, 4$ (they describe the deviation of DA's from its asymptotic form) for the P -wave production of pions are negative and $B_{41}^\parallel(w^2)$ increases (by absolute value) whereas $B_{21}^\parallel(w^2)$ decreases with increasing of w^2 .

Here we quote the result for the chirally even ρ meson DA extracted from $2\pi\text{DA}$ using eqs. (5.14,5.16):

$$\varphi_\rho^\parallel(z) = 6z(1-z)[1 - 0.14 C_2^{3/2}(2z-1) - 0.01 C_4^{3/2}(2z-1) + \dots], \quad (6.15)$$

with normalization constant $f_\rho^\parallel = 190 \text{ MeV}$ according to eq. (5.16) in a good agreement with experimental value $f_\rho^\parallel = 195 \pm 7 \text{ MeV}$ [31]. The ρ meson DA's were the subject of the QCD sum rules calculations [23, 32, 33], our result eq. (6.15) is in a qualitative disagreement with the results of QCD sum rule calculations, the sign of $a_2^\parallel$ obtained here is opposite to the QCD sum rules predictions $a_2^\parallel = 0.18 \pm 0.1$ [32] and $a_2^\parallel = 0.08 \pm 0.02$, $a_4^\parallel = -0.08 \pm 0.03$ [33] (these results refer to normalization point $\mu = 1 \text{ GeV}$). The difference in the sign of ours and QCD sum rule calculations can be traced back to the significantly different values of $a_2^{(\pi)}$. Indeed, using the eq. (5.14) and the crossing relation (4.8) we can relate $a_2^{(\rho)}$, $a_2^{(\pi)}$ and the third moment of the quark distribution in a pion in the following way:

$$\begin{aligned} a_2^{(\rho)} = B_{21}^{I=1}(0) \exp(c_1^{(21)} m_\rho^2) &= (a_2^{(\pi)} - B_{23}^{I=1}(0)) \exp(c_1^{(21)} m_\rho^2) \\ &= (a_2^{(\pi)} - \frac{7}{6} M_3^{(\pi)}) \exp(c_1^{(21)} m_\rho^2). \end{aligned} \quad (6.16)$$

The value of $a_2^{(\pi)}$ in the instanton model of the QCD vacuum is very small (see eq. (6.13) and ref. [28]), such that $a_2^{(\pi)} < \frac{7}{6} M_3^{(\pi)}$. On contrary the value of $a_2^{(\pi)}$ in the QCD sum rule calculations [34, 35] larger than $\frac{7}{6} M_3^{(\pi)}$, what implies the positive sign of $a_2^{(\rho)}$.

Using eq. (5.14) and the value of the subtraction constants from Table 1 we can make a rough estimate of the shape of the ρ' chirally even DA ($m_{\rho'} = 1.465 \text{ GeV}$):

$$\varphi_{\rho'}^\parallel(z) = 6z(1-z)[1 - 0.055 C_2^{3/2}(2z-1) - 0.064 C_4^{3/2}(2z-1) + \dots]. \quad (6.17)$$

This is just a rough estimate because the mass of ρ' is rather big and importance of neglected in the derivation of eq. (5.9) high energy states increases and hence the accuracy of the solution eq. (5.9) becomes poor. But even from this rough estimate we see an interesting tendency: the coefficients in front of the Gegenbauer polynomials with $n > 2$ in the expansion of ρ' meson chirally even DA become comparable with the leading nontrivial ($n = 2$) coefficient and the DA develops nodes.

Now we turn to results for the chirally odd 2π DA eq. (2.2). Simple calculations of matrix element eq. (2.3) give the following value of the constant $f_{2\pi}$

$$f_{2\pi} = \frac{8\pi^2 f_\pi^2}{N_c M} \approx 650 \text{ MeV} . \quad (6.18)$$

Calculations of $B_{nl}^\perp(w^2)$ in the low energy region are straightforward (details will be published elsewhere [36]), the results are:

$$B_{01}^\perp(w^2) = 1 + \frac{w^2}{12M^2} + \dots , \quad (6.19)$$

$$B_{21}^\perp(w^2) = \frac{7}{36} \left(1 - \frac{w^2}{30M^2} + \dots\right), \quad (6.20)$$

$$B_{23}^\perp(w^2) = \frac{7}{36} \left(1 + \frac{w^2}{30M^2} + \dots\right), \quad (6.21)$$

$$B_{41}^\perp(w^2) = \frac{11}{225} \left(1 - \frac{5w^2}{168M^2} + \dots\right), \quad (6.22)$$

$$B_{43}^\perp(w^2) = \frac{77}{675} \left(1 - \frac{w^2}{630M^2} + \dots\right), \quad (6.23)$$

$$B_{45}^\perp(w^2) = \frac{11}{135} \left(1 + \frac{w^2}{56M^2} + \dots\right) . \quad (6.24)$$

From these equations we get immediately the values of the subtraction constants $c_k^{\perp(nl)}$, they are summarized in Table 1. The value of “tensor” isovector radius of pion is from eq. (6.19) (without chiral loops!):

$$\langle r^2 \rangle_t = \frac{1}{2M^2} \approx 0.16 \text{ fm}^2 . \quad (6.25)$$

The chiral loops are the same as for charge radius.

Using the results from Table 1 for $c_k^{\perp(nl)}$ and eqs. (5.14,5.16) we obtain the chirally odd ρ -meson DA:

$$\varphi_\rho^\perp(z) = 6z(1-z)[1 + 0.11 C_2^{3/2}(2z-1) + 0.03 C_4^{3/2}(2z-1) + \dots] , \quad (6.26)$$

and the normalization constant $f_\rho^\perp$:

$$f_\rho^\perp = f_\rho^\parallel \frac{m_\rho}{f_{2\pi}} \exp[(\langle r^2 \rangle_t - \langle r^2 \rangle_{\text{e.m.}})m_\rho^2/6] \approx 140 \text{ MeV} . \quad (6.27)$$

[This value refers to the normalization point $\mu \sim 1/\bar{\rho} \approx 600$ MeV]. The QCD sum rule calculations [32, 33] give $f_\rho^\perp = 160 \pm 10$ MeV $a_2^\perp = 0.2 \pm 0.1$ [32] and $f_\rho^\perp = 169 \pm 5$ MeV $a_2^\perp = 0.36 \pm 0.03$ [33] at the normalization point $\mu = 1$ GeV in fair agreement with our results.

The results for parameters of the ρ meson DA's are summarized in Table 2 where we also quote the results of the QCD sum rule calculations [32, 33]. With the help of Table 1 and eqs. (5.9, 5.18) the reader can easily construct DA's of isovector mesons of any spin.

6.2 Note on isoscalar 2π DA's

The quark isoscalar chirally even 2π DA's under ERBL evolution [9] mix with gluon 2π DA's. In the instanton vacuum model considered here the gluon 2π DA's are parametrically small at low normalization point ($\mu \sim 1/\bar{\rho}$) in the instanton packing fraction (see ref. [38]) and hence are zero at the level of accuracy considered here.

The second moment of the chirally even 2π DA eq. (2.1) is related to the quark part of the (symmetric) energy-momentum tensor $T_q^{\mu\nu}$:

$$\int_0^1 dz (2z - 1) \Phi_{\parallel}^{I=0}(z, \zeta, w^2) = \frac{1}{2P^{+2}}{}_{out} \langle \pi^a(p_1) \pi^b(p_2) | T_q^{++} | 0 \rangle . \quad (6.28)$$

In the instanton model of the QCD vacuum (see [38]) the gluon part of the energy-momentum tensor (projected on the light-cone direction n^μ !) is parametrically small in the packing fraction of the instanton liquid. Therefore here we can use, the so-called, quark-antiquark approximation, *i.e.* $T_q^{++} = T^{++}$ + small corrections. At this level of accuracy we have (*cf.* equation (4.7)):

$$\int_0^1 dz (2z - 1) \Phi_{\parallel}^{I=0}(z, \zeta, w^2 = 0) = -2\zeta(1 - \zeta) . \quad (6.29)$$

The model calculations of the isoscalar chirally even 2π DA give the following result for its second moment:

$$\int_0^1 dz (2z - 1) \Phi_{\parallel}^{I=0}(z, \zeta, w^2) = -2\zeta(1 - \zeta) \left[1 + \frac{N_c w^2}{48\pi^2 f_\pi^2} + \dots \right] . \quad (6.30)$$

We see that the overall normalization is fixed by the energy momentum conservation and the radius of pion form factor of the energy momentum tensor is:

$$\langle r^2 \rangle_{EMT} = \frac{N_c}{8\pi^2 f_\pi^2} = 0.18 \text{ fm}^2 . \quad (6.31)$$

From eq. (6.30) we get immediately the values of the low-energy subtraction constant:

$$B_{10}^{\parallel}(0) = -\frac{5}{9}, \quad B_{12}^{\parallel}(0) = \frac{5}{9}, \quad (6.32)$$

$$\frac{d}{dw^2} \log B_{10}^{\parallel}(w^2)|_{w^2=0} = \frac{d}{dw^2} \log B_{12}^{\parallel}(w^2)|_{w^2=0} = \frac{N_c}{48\pi^2 f_\pi^2} = 0.73 \text{ GeV}^{-2}. \quad (6.33)$$

Obviously the soft pion theorem eq. (3.10) is satisfied. Let us use the above model calculations and crossing relation (4.2) to estimate the low momentum transfer behaviour of the second moments of skewed quark distribution in a pion. From eq. (4.2) for the second moment we have

$$\begin{aligned} \int_{-1}^1 dX X H^{I=0}(X, \xi, t) &= \frac{3}{5} B_{10}^{\parallel}(t) \xi^2 + \frac{3}{5} B_{12}^{\parallel}(t) \xi^2 C_2^{1/2} \left(\frac{1}{\xi} \right) \\ &= \frac{3}{5} [B_{10}^{\parallel}(t) \xi^2 + B_{12}^{\parallel}(t) \frac{3 - \xi^2}{2}] . \end{aligned} \quad (6.34)$$

Now if we substitute in the above equation the result of the model calculation (6.33) we get:

$$\int_{-1}^1 dX X H^{I=0}(X, \xi, t) = \frac{1}{2} (1 - \xi^2) \left[1 + \frac{N_c t}{48\pi^2 f_\pi^2} + \dots \right]. \quad (6.35)$$

7 Two-pion distribution amplitudes in hard pions production processes

The dependence of the two-pion hard production amplitude (1.3) on the $2\pi\text{DA}$'s factorizes into the factor:

$$A \propto \int_0^1 \frac{dz}{z(1-z)} \Phi_{\parallel}^{I=1}(z, \zeta, w^2; \bar{Q}^2), \quad (7.1)$$

for the two pions in the isovector state, and

$$A \propto \int_0^1 dz \frac{(2z-1)}{z(1-z)} \Phi_{\parallel}^{I=0}(z, \zeta, w^2; \bar{Q}^2), \quad (7.2)$$

for the pions in the isoscalar state (*e.g.* for the $\pi^0\pi^0$ production). In the above equations we showed also the dependence of the $2\pi\text{DA}$'s on the scale of the process $\bar{Q}^2$, which is governed by the ERBL evolution equation [9].

For the process (1.3) at small x_{Bj} (see *e.g.* recent measurements [39]) the production of two pions in the isoscalar channel is strongly suppressed relative to the isovector channel because the former is mediated by C -parity odd exchange. At asymptotically large Q^2 one can use the asymptotic form of isovector $2\pi\text{DA}$:

$$\lim_{\bar{Q}^2 \rightarrow \infty} \Phi_{\parallel}^{I=1}(z, \zeta, w^2; \bar{Q}^2) = 6F_\pi(w^2) z(1-z)(2\zeta-1), \quad (7.3)$$

where $F_\pi(w^2)$ is the pion e.m. form factor in the time-like region. Therefore the shape of $\pi^+\pi^-$ mass spectrum in the hard electroproduction process (1.3) at small x_{Bj} and

asymptotically large Q^2 should be determined completely by the pion e.m. form factor in time-like region:

$$\lim_{Q^2 \rightarrow \infty} A \propto e^{i\delta_1^1(w^2)} |F_\pi(w^2)| (2\zeta - 1). \quad (7.4)$$

Deviation of the $\pi^+\pi^-$ mass spectrum from its asymptotic form eq. (7.4) (“skewing”) can be parametrized at small x_{Bj} and large $\bar{Q}^2$ in the form:

$$\begin{aligned} A \propto e^{i\delta_1^1(w^2)} |F_\pi(w^2)| & \left[1 + B_{21}(0; \mu_0) \exp\{c_1^{(21)} w^2\} \left(\frac{\alpha_s(\bar{Q}^2)}{\alpha_s(\mu_0)} \right)^{50/(99-6N_f)} \right] (2\zeta - 1) + \\ & e^{i\delta_3^1(w^2)} B_{23}(0; \mu_0) \exp\{b_{23} w^2 + R_3^1(w^2)\} \left(\frac{\alpha_s(\bar{Q}^2)}{\alpha_s(\mu_0)} \right)^{50/(99-6N_f)} C_3^{1/2} (2\zeta - 1) \\ & + \text{higher powers of } 1/\log(\bar{Q}^2), \quad (7.5) \end{aligned}$$

here

$$R_l^I(w^2) = \text{Re} \frac{w^4}{\pi} \int_{4m_\pi^2}^\infty ds \frac{\delta_l^I(s)}{s^2(s - w^2 - i0)} \quad \text{and} \quad b_{nl} = \frac{d}{dw^2} \log B_{nl}(w^2) \Big|_{w^2=0}. \quad (7.6)$$

We see that the deviation of the $\pi\pi$ invariant mass spectrum from its asymptotic form eq. (7.4) in this approximation is characterized by a few low-energy constants ($B_{21}(0)$, $B_{23}(0)$, $c_1^{(21)}$, b_{23}), other quantities– the pion e.m. form factor and the $\pi\pi$ phase shifts–are known from low-energy experiments. In principle, using the parametrization (7.6) one can extract the values of these low-energy parameters from the shape of $\pi\pi$ spectrum (not absolute cross section!) in diffractive production experiments. Knowing them one can obtain the deviation of the π meson DA (see eq. (3.9))

$$a_2^{(\pi)} = B_{21}(0) + B_{23}(0),$$

and the ρ meson DA (see eq. (5.14))

$$a_2^{(\rho)} = B_{21}(0) \exp(c_1^{(21)} m_\rho^2),$$

from their asymptotic form $6z(1-z)$, as well as the normalization constants for the DA of isovector resonances of spin three (see eq. (5.19)). Additionally the parameter $B_{23}(0)$ is related by the crossing relations to the third moment of the valence quark distribution in a pion (see eq. (4.6)):

$$M_3^\pi = \frac{6}{7} B_{23}(0). \quad (7.7)$$

We see that, in principle, the diffractive production of two pions can be used to get an information about quark distribution in a pion.

In analysis of experiments on two pion diffractive production off nucleon (see e.g. [39]) the non-resonant background is described by Söding parametrization [41], which takes into account rescattering of produced pions on final nucleon. Let us note however that in a case of hard ($Q^2 \rightarrow \infty$) diffractive production the final state interaction of pions with residual nucleon is suppressed by powers of $1/Q^2$ relative to the leading twist amplitude. Here we proposed alternative leading-twist parametrization (7.6) describing the so-called “skewing” of two pion spectrum.

8 Conclusions

We showed that the two-pion distribution amplitudes are the most general object to describe the hard electroproduction of two pions. Using Watson final state interaction theorem and the soft pion theorems proved here, we can determine the 2π DA's in a wide region of pion invariant masses in term of the pion phase shifts and a few low-energy (subtraction) constants. The former are known from data on $\pi\pi$ scattering (soft physics), the latter are non-perturbative input which we estimated here using the instanton model of the QCD vacuum. We showed that these non-perturbative characteristics can, in principle, be extracted from the shape of $\pi\pi$ mass spectra in diffractive pions production experiments. Then they can be used to determine, say, the deviation of the π meson DA (see eq. (3.9)) and ρ meson DA (see eq. (5.14)) from their asymptotic form $6z(1-z)$, and hence to obtain non-perturbative information about structure of mesons. We demonstrated (see section 5.1) that DA's of resonances of any spin can be determined from 2π DA's.

We derive the crossing relation which relates various distribution amplitudes to the quark distributions in a pion.

The soft pion theorems considered in section 3 give an example of how low-energy (chiral) physics can be studied in hard processes (see recent work in this direction [40]).

The methods developed here can be easily generalized for more complicated cases. One of examples is generalized $N \rightarrow \pi N$ skewed parton distributions [12, 13] entering the QCD description of the “non-diagonal” deeply-virtual Compton scattering $\gamma_L^* + p \rightarrow \gamma + \pi N$.

9 Acknowledgments

I am grateful to P. Ball, V. Braun, L. Frankfurt, K. Goeke, P. Pobylitsa, A. Radyushkin, A. Schäfer, M. Strikman, O. Teryaev and C. Weiss for inspiring discussions. Special thanks to D.I. Diakonov for many valuable remarks. This work has been supported in part by the BMFB (Bonn), Deutsche Forschungsgemeinschaft (Bonn) and by COSY (Jülich).

A Appendix

Due to the kinematical constraint (2.5) the physical region of variable ζ is bounded to the interval

$\zeta_- \leq \zeta \leq \zeta_+$, where

$$\zeta_{\pm} = \frac{1}{2} \left(1 \pm \sqrt{1 - \frac{4m_{\pi}^2}{w^2}} \right). \quad (\text{A.1})$$

It implies that the Gegenbauer polynomials $C_l^{1/2}(2\zeta - 1)$ used in the expansion (2.9) are not orthogonal on the interval $\zeta_- \leq \zeta \leq \zeta_+$. The correct set of polynomials in this case is:

$$\mathcal{P}_l(\zeta) = C_l^{1/2} \left(\frac{2\zeta - 1}{\sqrt{1 - \frac{4m_{\pi}^2}{w^2}}} \right). \quad (\text{A.2})$$

It is easy to check that these polynomials are orthogonal on the interval $\zeta_- \leq \zeta \leq \zeta_+$:

$$\int_{\zeta_-}^{\zeta_+} d\zeta \mathcal{P}_l(\zeta) \mathcal{P}_k(\zeta) = \begin{cases} 0, & l \neq k, \\ \frac{1}{2l+1} \sqrt{1 - \frac{4m_{\pi}^2}{w^2}}, & l = k. \end{cases} \quad (\text{A.3})$$

In the case of w^2 close to the threshold $4m_{\pi}^2$ the expansion (2.9) is modified:

$$\Phi_{\parallel, \perp}(z, \zeta, w^2) = 6z(1-z) \sum_{n=0}^{\infty} \sum_{l=0}^{n+1} B_{nl}^{\parallel, \perp}(w^2) C_n^{3/2}(2z-1) \mathcal{P}_l(\zeta). \quad (\text{A.4})$$

From the normalization condition (2.8) we conclude that

$$B_{01}^{\parallel} = \sqrt{1 - \frac{4m_{\pi}^2}{w^2}} F_{\pi}^{\text{e.m.}}(w^2). \quad (\text{A.5})$$

Also the equations in section 5 are modified near threshold. The phase space volume is:

$$\int d(\text{phase space}) = \frac{1}{64\pi^2} \int_{\zeta_-}^{\zeta_+} d\zeta' \int_0^{2\pi} d\theta_{\perp}. \quad (\text{A.6})$$

The scattering angle in the c.m. system is expressed in terms light cone variables as following

$$\begin{aligned} \cos(\theta_{cm}) &= \frac{1}{1 - 4m_{\pi}^2/w^2} \left[1 - 2(\zeta + \zeta' - 2\zeta\zeta') \right. \\ &\quad \left. - 4\sqrt{(\zeta(1-\zeta) - m_{\pi}^2/w^2)(\zeta'(1-\zeta') - m_{\pi}^2/w^2)} \cos(\theta_{\perp}) \right]. \end{aligned} \quad (\text{A.7})$$

With this equality the expression (5.5) is modified as follows:

$$\int_0^{2\pi} d\theta_{\perp} P_l[\cos(\theta_{cm})] = 2\pi \mathcal{P}_l(\zeta) \mathcal{P}_l(\zeta') . \quad (\text{A.8})$$

This demonstrates that the set of polynomials (A.2) corresponds to the expansion in partial waves of produced pions.

References

- [1] J. Collins, L. Frankfurt, and M. Strikman, Phys. Rev. **D56** (1997) 2982.
- [2] A. V. Radyushkin, Phys. Rev. **D56** (1997) 5524
- [3] X. Ji, Phys. Rev. **D55** (1997) 7114.
- [4] F.M. Dittes, D. Müller, D. Robaschik, B. Geyer, J. Horejsi Phys.Lett. **B209** (1988) 325.
- [5] D. Müller, D. Robaschik, B. Geyer, F.M. Dittes, J. Horejsi, Fortschr. Phys. **42** (1994) 101.
- [6] X. Ji, J. Phys. **G24** (1998) 1181.
- [7] M. Diehl, T. Gousset, B. Pire and O. Teryaev, Phys. Rev. Lett. **81** (1998) 1782.
- [8] M.V. Polyakov and C. Weiss, Bochum University preprint RUB-TPII-7/98, hep-ph/9806390, to appear in Phys. Rev. D.
- [9] G.P. Lepage and S.J. Brodsky, Phys. Lett. **B 87** (1979) 359; A.V. Efremov and A.V. Radyushkin, Phys. Lett. **B 94** (1980) 245.
- [10] M.A. Shifman and M.I. Vysotsky, Nucl. Phys. **B 186**, 475 (1981).
- [11] O. Teryaev, talk given at “XIV International Seminar on High Energy Physics Problems”, August 1998.
- [12] L.L. Frankfurt, M.V. Polyakov, M. Strikman, hep-ph/9808449.
- [13] M.V. Polyakov, talk given at “Baryon98”, September 1998, to appear in proceedings.
- [14] M.V. Polyakov, talk given at the conference “Particle and Nuclear Physics with CEBAF at Jefferson Lab”, Dubrovnik, 3-10 November 1998, to be published in the proceedings, hep-ph/9901315.
- [15] M.V. Polyakov and C. Weiss, Bochum University preprint, RUB-TP2-01/99, hep-ph/9902451.
- [16] A.V. Radyushkin, Phys.Rev. **D59** (1999) 014030; A.V. Radyushkin, hep-ph/9810466.

- [17] K.M. Watson, Phys. Rev. **95** (1955) 228.
- [18] F. Guerrero and A. Pich, Phys. Lett. **B412** (1997) 382.
- [19] R. Omnes, Nuovo Cim. **8** (1958) 316.
- [20] F. Guerrero, Phys. Rev. **D57** (1998) 4136.
- [21] F. Guerrero, J.A. Oller, Nucl. Phys. **B537** (1999) 459.
- [22] D. Diakonov and V. Petrov, Nucl. Phys. **B 272** (1986) 457; LNPI preprint LNPI-1153 (1986), published (in Russian) in: Hadron matter under extreme conditions, Naukova Dumka, Kiev (1986), p.192.
- [23] For a review, see V.L. Chernyak and A.R. Zhitnitsky, Phys. Rept. **112** (1984) 173.
- [24] J. Bartels and M. Loewe, Z. Phys. **C12** (1982) 283;
M. G. Ryskin, Z. Phys. **C57** (1993) 89;
S. J. Brodsky *et al.* Phys. Rev. **D50** (1994) 3134;
L. Frankfurt, W. Koepf and M. Strikman, Phys. Rev. **D54** (1996) 3194.;
L. Mankiewicz, G. Piller and T. Weigl, Eur. Phys. J. **C5** (1998) 119.
- [25] A. Ali, V. M. Braun and H. Simma, Z. Phys. **C63** (1994) 437.
- [26] D.I.Diakonov, Yu.V.Petrov and P.V.Pobylitsa, Nucl. Phys. **B306** (1988) 809
- [27] V.Yu. Petrov and P.V. Pobylitsa, hep-ph/9712203.
- [28] V.Yu. Petrov *et al.*, Bochum University preprint RUB-TPII-8/98, hep-ph/9807229, to appear in Phys. Rev. **D**.
- [29] S. R. Amedolia *et al.*, Nucl. Phys. **B277** (1986) 168.
- [30] J. Gasser, H. Leutwyller, Nucl. Phys. **B250** (1985) 465.
- [31] Particle Data Group, Phys. Rev. **D54** (1996), 1.
- [32] P. Ball and V. M. Braun, Phys. Rev. **D54** (1996) 2186.
- [33] A.P. Bakulev and S.V. Mikhailov, Phys. Lett. **B436** (1998) 351.
- [34] V. Braun and I. Filyanov, Z. Phys. **C44** (1989) 157.
- [35] S. V. Mikhailov and A.V. Radyushkin, Phys. Rev. **D45** (1992) 1754.
- [36] M.V. Polyakov *et al.*, in preparation.
- [37] V.Yu. Petrov, P.V. Pobylitsa, M.V. Polyakov, I. Börnig, K. Goeke, and C. Weiss, Phys. Rev. **D 57** (1998) 4325.
- [38] D.I. Diakonov, M.V. Polyakov and C. Weiss, Nucl. Phys. **B 461**, 539 (1996);
J. Balla, M.V. Polyakov and C. Weiss, Nucl. Phys. **B 510**, 327 (1997).

- [39] ZEUS Collaboration, hep-ex/9808020.
- [40] M. Eides, L. Frankfurt and M. Strikman, hep-ph/9809277
- [41] P. Söding, Phys. Lett. **B19** (1966) 702.

(nl)	$B_{nl}^{\parallel}(0)$	$B_{nl}^{\perp}(0)$	$c_1^{\parallel(nl)} \text{ GeV}^{-2}$	$c_1^{\perp(nl)} \text{ GeV}^{-2}$	$\frac{d}{dw^2} \log B_{nl}^{\parallel}(0) \text{ GeV}^{-2}$	$\frac{d}{dw^2} \log B_{nl}^{\perp}(0) \text{ GeV}^{-2}$
(01)	1	1	0	0	1.46	0.68
(21)	-0.2	0.2	-0.6	-0.95	0.85	-0.27
(23)	0.26	0.2	0	0	0.65	0.27
(41)	-0.006	0.05	1.1	-0.92	2.5	-0.24
(43)	-0.12	0.11	-0.36	-0.28	0.30	-0.01
(45)	0.14	0.08	0	0	0.36	0.15

Table 1: Values of the subtraction constants entering eqs. (5.9,5.14) calculated in the model of instanton vacuum. Isovector case.

	our results	Ball & Braun	Bakulev & Mikhailov
$f_{\rho}^{\parallel} \text{ (MeV)}$	190	198 ± 7	201 ± 5
$f_{\rho}^{\perp} \text{ (MeV)}$	140	160 ± 10	169 ± 5
$a_2^{\parallel}$	-0.14	0.18 ± 0.10	0.08 ± 0.02
$a_4^{\parallel}$	-0.01	—	-0.08 ± 0.03
$a_2^{\perp}$	0.11	0.2 ± 0.10	0.36 ± 0.03
$a_4^{\perp}$	0.03	—	—

Table 2: Parameters of the twist-2 vector meson distribution amplitudes extracted from the 2π DA's compared with QCD sum rule calculations [34,35]. Our results refer to normalization point $\mu \approx 600 \text{ MeV}$, QCD sum rule calculations to $\mu \approx 1 \text{ GeV}$.




Figure 1: The two-pion chirally even isovector distribution amplitude $\Phi_{\parallel}^{I=1}(z, \zeta, w^2 = 0)$ at various values of ζ . *Solid line:* $\zeta = 0.1$. *Dashed line:* $\zeta = 0.3$. *Dotted line:* $\zeta = 0.45$